

LATIN k-CUBES

JOSEPH ARKIN

197 Old Nyack Turnpike, Spring Valley, New York 10977

and

E. G. STRAUS

University of California, Los Angeles 90024

1. LATIN SQUARES

A Latin square of order n is an $n \times n$ square in which each of the numbers $0, 1, \dots, n-1$ occurs exactly once in each row and exactly once in each column. For example

$$\begin{array}{ccc}
 0 & 1 & 0 & 1 & 2 & 0 & 1 & 2 & 3 \\
 1 & 0 & 1 & 2 & 0 & 1 & 2 & 3 & 0 \\
 & & 2 & 0 & 1 & 2 & 3 & 0 & 1 \\
 & & & & & 3 & 0 & 1 & 2
 \end{array}$$

are Latin squares of order $2, 3, 4$, respectively. Two Latin squares of order n are orthogonal, if when one is superimposed on the other, every ordered pair $00, 01, \dots, n-1, n-1$ occurs. Thus

$$\begin{array}{ccc}
 0 & 1 & 2 & & 0 & 1 & 2 & & 0 & 0 & 1 & 1 & 2 & 2 \\
 1 & 2 & 0 & \text{and} & 2 & 0 & 1 & \text{superimpose to} & 1 & 2 & 2 & 0 & 0 & 1 \\
 2 & 0 & 1 & & 1 & 2 & 0 & & 2 & 1 & 0 & 2 & 1 & 0
 \end{array}$$

and therefore are orthogonal squares of order 3. A set of Latin squares of order n is orthogonal if every two of them are orthogonal. As an example the 4×4 square of triples

$$\begin{array}{cccc}
 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 & 3 & 3 & 3 \\
 1 & 2 & 3 & 0 & 3 & 2 & 3 & 0 & 1 & 2 & 1 & 0 \\
 2 & 3 & 1 & 3 & 2 & 0 & 0 & 1 & 3 & 1 & 0 & 2 \\
 3 & 1 & 2 & 2 & 0 & 3 & 1 & 3 & 0 & 0 & 2 & 1
 \end{array}$$

represents three mutually orthogonal squares of order 4 since each of the 16 pairs $00, 01, \dots, 33$ occurs in each of the three possible positions among the 16 triples.

There cannot exist more than $n-1$ mutually orthogonal Latin squares of order n , and the existence of such a complete system is equivalent to the existence of a finite projective plane of order n , that is a system of n^2+n+1 points and n^2+n+1 lines with $n+1$ points on each line. If n is a power of a prime there exist finite fields of order n which can be used to construct finite projective planes of order n . So, for $n=2, 3, 4, 5, 7, 8, 9$ there exist complete systems of $n-1$ orthogonal Latin squares of order n . We have listed the examples $n=2, 3, 4$, above. It is known [2] that there are no orthogonal Latin squares of order 6 and that there are at least two orthogonal Latin squares of every order $n > 2, n \neq 6$. In fact, the number of mutually orthogonal Latin squares of order n goes to infinity with n [3]. However no case of a complete system of $n-1$ orthogonal Latin squares is known for any n which is not a power of a prime.

2. LATIN CUBES

We can generalize all these concepts to $n \times n \times n$ cubes and cubes of higher dimensions.

A Latin cube of order n is an $n \times n \times n$ cube (n rows, n columns and n files) in which the numbers $0, 1, \dots, n-1$ are entered so that each number occurs exactly once in each row, column and file. If we list the cube in terms of the n squares of order n which form its different levels we can list the cubes

$$\begin{array}{cc}
 0 & 1 & & 1 & 0 & & \text{and} & 0 & 1 & 2 & & 1 & 2 & 0 & & 2 & 0 & 1 \\
 1 & 0 & & 0 & 1 & & & 1 & 2 & 0 & & 2 & 0 & 1 & & 0 & 1 & 2 \\
 & & & & & & & 2 & 0 & 1 & & 0 & 1 & 2 & & 1 & 2 & 0
 \end{array}$$

as Latin cubes of order two and three, respectively. Since even this method of listing becomes unwieldy for higher dimensions we also use a listing by indices. Thus we write the first cube as $A = (a_{ijk})$ with $a_{000} = 1, a_{010} = 1, a_{011} = 0, a_{100} = 1, a_{101} = 0, a_{110} = 0, a_{111} = 1$. In a similar manner we can describe four-dimensional cubes $A = (a_{ijk\ell})$ or order n , where each of the indices, i, j, k, ℓ ranges from 1 to n . Generally we can discuss k -cubes $A = (a_{i_1 i_2 \dots i_k})$ with k indices ranging from 1 to n . These cubes will be Latin k -cubes of order n if each of the n^k entries $a_{i_1 \dots i_k}$ is one of the numbers $0, 1, \dots, n-1$ so that $a_{i_1 \dots i_k}$ ranges over all these numbers as one of the indices varies from 1 to n while the other indices remain fixed.

Orthogonality of Latin cubes is now a relation among three cubes, or in general among k Latin k -cubes. That is, three Latin cubes of order n are orthogonal if, when superimposed, each ordered triple will occur. Similarly k Latin k -cubes are orthogonal if, when superimposed, each ordered k -tuple will occur. A set of at least k Latin k -cubes is orthogonal if every k of its cubes are orthogonal.

Theorem. If there exist two orthogonal Latin squares of order n then there exist 4 orthogonal Latin cubes of order n and k orthogonal Latin k -cubes for each $k > 3$.

Proof. Let $A = (a_{ij}), B = (b_{ij})$ be orthogonal Latin squares of order n . Define 4 cubes C, D, E, F of order n by

$$c_{ijk} = a_{a_{ij}, k}, d_{ijk} = a_{b_{ij}, k}, c_{ijk} = b_{a_{ij}, k}, f_{ijk} = b_{b_{ij}, k}, i, j, k = 0, 1, \dots, n-1.$$

Note that the squares A, B are used both as entries and as indices in the construction of the cubes. For example the pair of 3×3 Latin squares

$$\begin{array}{ccc}
 0 & 0 & 1 & 1 & 2 & 2 \\
 1 & 2 & 2 & 0 & 0 & 1 \\
 2 & 1 & 0 & 2 & 1 & 0
 \end{array}$$

leads to the four $3 \times 3 \times 3$ cubes

C:	0 1 2	1 2 0	2 0 1
	1 2 0	2 0 1	0 1 2
	2 0 1	0 1 2	1 2 0
D:	0 1 2	1 2 0	2 0 1
	2 0 1	0 1 2	1 2 0
	1 2 0	2 0 1	0 1 2
E:	0 2 1	1 0 2	2 1 0
	2 1 0	0 2 1	1 0 2
	1 0 2	2 1 0	0 2 1
F:	0 2 1	1 0 2	2 1 0
	1 0 2	2 1 0	0 2 1
	2 1 0	0 2 1	1 0 2

Superimposed these lead to a cube of quadruples

CDEF:	0000	1122	2211	1111	2200	0022	2222	0011	1100
	1221	2010	0102	2002	0121	1210	0110	1202	2021
	2112	0201	1020	0220	1012	2101	1001	2120	0212

where each ordered triple occurs in every one of the four possible positions in the quadruples. It is easy to see that C, D, E, F are Latin cubes. For example, for fixed ij the values $c_{ijk} = a_{a_{ij}, k}$ go through the a_{ij}^{th} row of A , that is, through the values $0, 1, \dots, n-1$. For fixed i, k the index a_{ij} goes through all the values in the i^{th} row of A , that is, through all values $0, 1, \dots, n-1$ and hence c_{ijk} goes through all values in the k^{th} column of A . Finally for fixed j, k the index a_{ij} goes through all values in the j^{th} column of A and therefore c_{ijk} again goes through all values in the k^{th} column of A .

To prove the orthogonality of, say, C, D, E we have to prove that for every triple (x, y, z) from $\{0, 1, \dots, n-1\}$ the equations

$$c_{ijk} = x, \quad d_{ijk} = y, \quad e_{ijk} = z$$

have a solution i, j, k . By the orthogonality of A and B the pair (x, z) occurs exactly once in the superimposed square AB so that the equations $a_{\varrho, k} = x, b_{\varrho, k} = z$ determine k and ϱ . Thus the equations

$$c_{ijk} = a_{a_{ij}, k} = x \quad \text{and} \quad e_{ijk} = b_{a_{ij}, k} = z$$

determine a_{ij} and k . Now, since A is a Latin square, there is exactly one occurrence of y in the k^{th} column of A so the equation

$$d_{ijk} = a_{b_{ij}, k} = y$$

determines b_{ij} and the pair (a_{ij}, b_{ij}) determines i, j ;

Thus for every triple (x, y, z) there is a unique triple (i, j, k) .

This construction is essentially that given by Arkin for 4 orthogonal $10 \times 10 \times 10$ cubes [1].

To prove the last part of the theorem we proceed by induction on k . Let A^1, \dots, A^k be orthogonal Latin k -cubes of order n , and write the entries of A^j as a_{i_1, \dots, i_k}^j . We now define $k+1$ orthogonal Latin $(k+1)$ -cubes B^1, \dots, B^{k+1} by

$$\begin{aligned} b_{i_1, \dots, i_{k+1}}^1 &= a_{i_1, \dots, i_k, i_{k+1}}^1 \\ &\vdots \\ b_{i_1, \dots, i_{k+1}}^k &= a_{i_1, \dots, i_k, i_{k+1}}^k \\ b_{i_1, \dots, i_{k+1}}^{k+1} &= b_{i_1, \dots, i_k, i_{k+1}}^1 \end{aligned}$$

We omit the proof that the B^j are Latin cubes, which is the same as before. In order to prove orthogonality we have to solve

$$B_{i_1 \dots i_{k+1}}^j = x_j \quad j = 1, \dots, k+1.$$

For any $(k+1)$ -tuple (x_1, \dots, x_{k+1}) from $\{0, 1, \dots, n-1\}$. Now, by the orthogonality of A and B the two equations

$$A_{a_{i_1, \dots, i_k}^1, i_{k+1}}^1 = x_1, \quad B_{a_{i_1, \dots, i_k}^1, i_{k+1}}^1 = x_{k+1}$$

determine $a_{i_1 \dots i_k}^1$ and i_{k+1} . Once i_{k+1} is determined the equations

$$A_{a_{i_1, \dots, i_k}^j, i_{k+1}}^j = x_j \quad j = 2, \dots, k$$

determine

$$a_{i_1 \dots i_k}^j \quad (j = 2, \dots, k).$$

Once the elements

$$a_{i_1 \dots i_k}^j \quad (j = 1, \dots, k)$$

are determined it follows from the orthogonality of the k -cubes A^1, \dots, A^k that the indices i_1, \dots, i_k are determined. Thus for every $(k+1)$ -tuple (x_1, \dots, x_{k+1}) there is a unique $(k+1)$ -tuple (i_1, \dots, i_{k+1}) with

$$B_{i_1 \dots i_{k+1}}^j = x_j \quad j = 1, \dots, k+1.$$

Since, as we mentioned, there are orthogonal Latin squares of every order $n \geq 2, n \neq 6$ we have the following.

Corollary. There exist orthogonal k -tuples of Latin k -cubes of order n for every $n > 2, n \neq 6$.

3. FINITE FIELDS

A field is a system of elements closed under the rational operations of addition, subtraction, multiplication and division (except by 0) subject to the usual commutative, associative and distributive laws. There exist finite fields with n elements if and only if n is a power of a prime p . The prime p is the characteristic of the field and we have $p\alpha = 0$ for every α in the field. Following are the addition and multiplication tables for the fields with 3 and 4 elements:

F_3	+	0	1	2	0	×	0	1	2
	0	0	1	2	0	0	0	0	0
	1	1	2	0	1	0	1	2	2
	2	2	0	1	2	0	2	1	1

	+	0	1	a	1+a		×	0	1	a	1+a
F_4	0	0	1	a	1+a	0	0	0	0	0	0
	1	1	0	1+a	a	1	0	1	a	1+a	
	a	a	1+a	0	1	a	0	a	1+a	1	
	1+a	1+a	a	1	0	1+a	0	1+a	1	a	

If there is a field F_n with n elements, that is if n is a power of a prime, we use the elements $\{f_1, f_2, \dots, f_n\}$ of F_n as indices to construct Latin squares, cubes, etc. We give the construction for cubes, but the generalization to k -cubes is easily seen.

Let α, β, γ be three nonzero elements of F_n then we can define the Latin cube $A = (a_{ijk})$ by

$$a_{ijk} = \alpha f_i + \beta f_j + \gamma f_k.$$

To see that A is a Latin cube consider, say, fixed i, j and see that γf_k runs through all elements of F_n as f_k does. Hence a_{ijk} runs through F_n as $k = 1, \dots, n$.

Now let $(\alpha, \beta, \gamma), (\alpha', \beta', \gamma')$ and $(\alpha'', \beta'', \gamma'')$ be three triples of nonzero elements of F_n so that the determinant

$$\begin{vmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{vmatrix} \neq 0.$$

Then the three Latin cubes

$$A = (a_{ijk}), \quad A' = (a'_{ijk}), \quad A'' = (a''_{ijk})$$

with

$$a_{ijk} = \alpha f_i + \beta f_j + \gamma f_k, \quad a'_{ijk} = \alpha' f_i + \beta' f_j + \gamma' f_k, \quad a''_{ijk} = \alpha'' f_i + \beta'' f_j + \gamma'' f_k$$

are orthogonal. This follows from the fact that for any triple (x, y, z) from F_n the three equations

$$a_{ijk} = x, \quad a'_{ijk} = y, \quad a''_{ijk} = z$$

have a unique solution f_i, f_j, f_k .

Now the Vandermonde determinants

$$\begin{vmatrix} 1 & a & a^2 \\ 1 & \beta & \beta^2 \\ 1 & \gamma & \gamma^2 \end{vmatrix} = (\beta - a)(\gamma - a)(\gamma - \beta)$$

are different from zero for any three distinct elements α, β, γ of F_n . Thus, letting a run through the nonzero elements of F_n we get $n - 1$ orthogonal Latin cubes of order n ,

$$A^a = (a^a_{ijk}), \quad a^a_{ijk} = f_i + a f_j + a^2 f_k.$$

The construction for a system of $n - 1$ orthogonal Latin k -cubes of order n proceeds in exactly the same way if we set

$$A^a = (a^a_{i_1 \dots i_k}), \quad a^a_{i_1 \dots i_k} = f_{i_1} + a f_{i_2} + \dots + a^{k-1} f_{i_k}$$

where a runs through the nonzero elements of F_n .

Theorem. If n is a power of a prime and $k < n$, then there exists a system of $n - 1$ orthogonal k -cubes of order n .

Our previous examples constructing four orthogonal cubes of orders 3 and 4 show that $n - 1$ is not necessarily the maximal number of orthogonal k -cubes of order n for $k > 2$. However, the orthogonal cubes constructed with the aid of finite fields satisfy additional properties. For each fixed value of k the squares

$$A^a_{i \cdot \cdot k} = (a^a_{ijk}) \quad i, j = 1, 2, \dots, n$$

form a complete system of $n - 1$ orthogonal Latin squares as a ranges through the nonzero elements of F_n , and similarly for each fixed i the squares

$$A^a_{i \cdot \cdot k} = (a^a_{ijk}) \quad j, k = 1, 2, \dots, n$$

form a complete system of orthogonal Latin squares. If n is a power of 2 then the third family of cross-sections

$$A^a_{i \cdot j \cdot} = (a^a_{ijk}) \quad i, k = 1, 2, \dots, n$$

form a complete system of orthogonal Latin squares for each fixed j , while for n odd we get a system of $(n - 1)/2$ orthogonal Latin squares, each square occurring twice.

Theorem. If n is a power of 2 then there exist $n - 1$ orthogonal Latin cubes of order n with the property that the corresponding plane sections form systems of $n - 1$ orthogonal Latin squares.

If n is a power of an odd prime then there exist $n - 1$ orthogonal Latin cubes with the property that the corresponding plane cross-sections in two directions form complete systems of orthogonal Latin squares, while the plane cross-sections in the third direction form a system of $(n - 1)/2$ orthogonal Latin squares, each square occurring twice.

Finally we observe that if we have orthogonal k -cubes of orders m and n then we can form their Kronecker products to obtain orthogonal k -cubes of order mn . That is from orthogonal k -cubes

$$A^1 = (a_{i_1 \dots i_k}^1), \dots, A^q = (a_{i_1 \dots i_k}^q); \quad B^1 = (b_{i_1 \dots i_k}^1), \dots, \quad B^q = (b_{i_1 \dots i_k}^q),$$

where the a 's run from 1 to m and the b 's from 1 to n we can form the orthogonal k -cubes C^1, \dots, C^q , where

$$C^j = (c_{i_1 \dots i_k}^j) \quad \text{and} \quad c_{i_1 \dots i_k}^j = (a_{i_1 \dots i_k}^j, b_{i_1 \dots i_k}^j)$$

so that the c 's run through all ordered pairs $(1, 1), \dots, (m, n)$ as the pairs $(i_1, j_1), \dots, (i_k, j_k)$ run through these ordered pairs. Thus we have the following.

Corollary. If

$$n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_s^{\alpha_s} \quad \text{and} \quad q = \min_{1 \leq j \leq s} p_j^{\alpha_j}$$

then for any $k < q$ there exist at least $q - 1$ orthogonal Latin k -cubes of order n .

The relation to finite k -dimensional projective spaces is not as immediate as it is for Latin squares, and we shall not discuss it here.

REFERENCES

1. J. Arkin, "A Solution to the Classical Problem of Finding Systems of Three Mutually Orthogonal Numbers in a Cube Formed by Three Superimposed $10 \times 10 \times 10$ Cubes," *The Fibonacci Quarterly*, Vol. 12, No. 2 (April 1974) pp. 133-140. Also, *Sugaku Seminar*, 13 (1974), pp. 90-94.
2. R.C. Bose, S.S. Shrikhande, and E.T. Parker, "Further Results on the Construction of Mutually Orthogonal Latin-Squares and the Falsity of Euler's Conjecture," *Canadian J. Math.*, 12 (1960), pp. 189-203.
3. S. Chowla, P. Erdős and E.G. Straus, "On the Maximal Number of Pair-wise Orthogonal Latin Squares of a Given Order," *Canadian J. Math.*, 12 (1960), pp. 204-208.
4. J. Arkin and V.E. Hoggatt, Jr., "The Arkin-Hoggatt Game and the Solution of a Classical Problem," *Jour. Recreational Math.*, Vol. 6, No. 2 (Spring, 1973), pp. 120-122.

This research was supported in part by NSF Grant No. GP-28696.

ON EXTENDING THE FIBONACCI NUMBERS TO THE NEGATIVE INTEGERS

M. G. MONZINGO

Southern Methodist University, Dallas, Texas 75275

A sequence of positive integers defined by the formula

$$(1) \quad x_{n+1} = ax_n + bx_{n-1}, \quad n \text{ a positive integer,}$$

is said to be extendable to the negative integers if (1) holds for n any integer. See page 28 of [1]. The purpose of this note is to show that the Fibonacci numbers form a sequence which is extendable to the negative integers in a unique way. In this note only nontrivial integral sequences will be considered.

[Continued on Page 308.]

THE FIBONACCI QUARTERLY

THE OFFICIAL JOURNAL OF
THE FIBONACCI ASSOCIATION



VOLUME 12

NUMBER 3

CONTENTS

Triangular Numbers	<i>V.E. Hoggatt, Jr., and Marjorie Bicknell</i>	221
On Fern's Theorem on the Expansion of Fibonacci and Lucas Numbers	<i>A.J.W. Hilton</i>	231
Argand Diagrams of Extended Fibonacci and Lucas Numbers	<i>F.J. Wunderlich, D.E. Shaw, and M.J. Hones</i>	233
A Pentagonal Arch	<i>Duane W. DeTemple</i>	235
A Generalization of the Hilton-Fern Theorem on the Expansion of Fibonacci and Lucas Numbers	<i>A.G. Shannon</i>	237
Some Aspects of Generalized Fibonacci Numbers	<i>J.E. Walton and A.F. Horadam</i>	241
An Extension of Fibonacci's Sequence	<i>P.J. deBruijn</i>	251
Spanning Trees and Fibonacci and Lucas Numbers	<i>A.J.W. Hilton</i>	259
On Polynomials Related to Tchebichef Polynomials of the Second Kind	<i>D. V. Jaiswal</i>	263
Corrigendum To: Enumeration of Two-Line Arrays	<i>L. Carlitz and Margaret Hodel</i>	266
Oresme Numbers	<i>A.F. Horadam</i>	267
Incredible Identities	<i>Daniel Shanks</i>	271
Some Further Identities for the Generalized Fibonacci Sequence $\{H_n\}$	<i>J.E. Walton and A.F. Horadam</i>	272
Explicit Expressions for Powers of Linear Recursive Sequences	<i>A.G. Shannon</i>	281
Latin k -Cubes	<i>Joseph Arkin and E.G. Straus</i>	288
On Extending the Fibonacci Numbers to the Negative Integers	<i>M.G. Monzingo</i>	292
A Method of Carlitz Applied to the K^{TH} Power Generating Function for Fibonacci Numbers	<i>A.G. Shannon</i>	293
A Constructed Solution of $\sigma(n) = \sigma(n + 1)$	<i>Richard Guy and Daniel Shanks</i>	299
The Design of the Four Binomial Identities: Moriarty Intervenes	<i>H.W. Gould</i>	300
Advanced Problems and Solutions	<i>Edited by Raymond E. Whitney</i>	309
Elementary Problems and Solutions	<i>Edited by A.P. Hillman</i>	313
OCTOBER		1974